

# Literatur

- [1] Araújo MG, Silva CO, Misawa M et al.: Alveolar socket healing: what can we learn? *Periodontol* 2000, 2015; 68(1): 122–34.
- [2] Parzeller M, Wenk M, Zedler B et al.: Aufklärung und Einwilligung bei ärztlichen Eingriffen. *Dtsch Arztebl International* 2007; 104(9): A–576EP.
- [3] Hugoson A, Kugelberg CF: The prevalence of third molars in a Swedish population. An epidemiological study. *Community Dent Health* 1988; 5(2): 121–38.
- [4] Artun J, Behbehani F, Thalib L: Prediction of maxillary third molar impaction in adolescent orthodontic patients. *Angle Orthod* 2005; 75(6): 904–11.
- [5] Fernandes MJ, Ogden GR, Pitts NB et al.: Incidence of symptoms in previously symptom-free impacted lower third molars assessed in general dental practice. *Nature Publishing Group* 2009; 207(5), E10–discussion: 218–9.
- [6] Chuang SK, Perrott DH, Susarla SM et al.: Age as a risk factor for third molar surgery complications. *YJOMS* 2007; 65(9): 1685–92.
- [7] Ades AG, Joondeph DR, Little RM et al.: A long-term study of the relationship of third molars to changes in the mandibular dental arch. *Am J Orthod Dentofacial Orthop* 1990; 97(4): 323–35.
- [8] Coulthard P, Bailey E, Esposito M et al.: Surgical techniques for the removal of mandibular wisdom teeth. *Cochrane Database Syst Rev* 2014; (7): CD004345.
- [9] Tolstunov L.: Influence of immediate post-extraction socket irrigation on development of alveolar osteitis after mandibular third molar removal: a prospective split-mouth study, preliminary report. *Nature Publishing Group* 2012; 213(12): 597–601.
- [10] Barone A, Marconcini S, Giacomelli L et al.: A randomized clinical evaluation of ultrasound bone surgery versus traditional rotary instruments in lower third molar extraction. *J Oral Maxillofac Surg* 2010; 68(2): 330–6.
- [11] Sivolella S, Berengo M, Bressan E et al.: Osteotomy for lower third molar germectomy: randomized prospective crossover clinical study comparing piezosurgery and conventional rotatory osteotomy. *J Oral Maxillofac Surg* 2011; 69(6): e15–23.
- [12] Lodi G, Figini L, Sardella A et al.: Antibiotics to prevent complications following tooth extractions. *Cochrane Database Syst Rev* 2012; 11: CD003811.
- [13] Marcussen KB, Laulund AS, Jørgensen HL et al.: A Systematic Review on Effect of Single-Dose Preoperative Antibiotics at Surgical Osteotomy Extraction of Lower Third Molars. *J Oral Maxillofac Surg* 2016; 74(4): 693–703.
- [14] Schultze-Mosgau S, Schmelzeisen R, Frölich JC et al.: Use of ibuprofen and methylprednisolone for the prevention of pain and swelling after removal of impacted third molars. *YJOMS* 1995; 53(1): 2–7–discussion7–8.
- [15] Freda NM, Keenan AV.: Moderate evidence to recommend submucosal injection of dexamethasone in reducing post-operative oedema and pain after third molar extraction. *Evid Based Dent* 2016; 17(2): 58–9.
- [16] Noguera-Gonzalez D, Cerda-Cristerna BI, Chavarria-Bolaños D et al.: Efficacy of preoperative ibuprofen on the success of inferior alveolar nerve block in patients with symptomatic irreversible pulpitis: a randomized clinical trial. *Int Endod J* 2013; 46(11): 1056–62.
- [17] Kunkel M: S2k-Leitlinie - Operative Entfernung von Weisheitszähnen [Internet]. AWMF-Register Nr. 007/003 Klasse: S2k. [cited 2017 Jul 4]. [www.awmf.org/uploads/tx\\_szleitlinien/007-003l\\_S2k\\_Weisheitszahnentfernung\\_01-2013.pdf](http://www.awmf.org/uploads/tx_szleitlinien/007-003l_S2k_Weisheitszahnentfernung_01-2013.pdf)
- [18] Blakey GH, Marciani RD, Haug RH et al.: Periodontal pathology associated with asymptomatic third molars. *YJOMS* 2002; 60(11): 1227–33.
- [19] Yildirim G, Ataoğlu H, Mihmanlı A et al.: Pathologic changes in soft tissues associated with asymptomatic impacted third molars. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2008; 106(1): 14–8.
- [20] Yamaoka M, Furusawa K, Ikeda M et al.: Root resorption of mandibular second molar teeth associated with the presence of the third molars. *Aust Dent J* 1999; 44(2): 112–6.

- [21] Fisher EL, Moss KL, Offenbacher S et al.: Third molar caries experience in middle-aged and older Americans: a prevalence study. *J Oral Maxillofac Surg* 2010; 68(3): 634–40.
- [22] Rakprasitkul S.: Pathologic changes in the pericoronal tissues of unerupted third molars. *Quintessence Int* 2001; 32(8): 633–8.
- [23] Bezerra TP, Studart-Soares EC, Pita-Neto IC et al.: Do third molars weaken the mandibular angle? *Med Oral Patol Oral Cir Bucal* 2011; 16(5): e657–63.
- [24] Güllicher D, Gerlach KL.: Inzidenz, Risikofaktoren und Verlauf von Sensibilitätsstörungen nach operativer Weisheitszahnentfernung. *Mund* 2000; 4(2): 99–104.
- [25] Libersa P, Roze D, Cachart T et al.: Immediate and late mandibular fractures after third molar removal. *YJOMS* 2002; 60(2): 163–5–discussion165–6.
- [26] Perry PA, Goldberg MH.: Late mandibular fracture after third molar surgery: a survey of Connecticut oral and maxillofacial surgeons. *YJOMS* 2000; 58(8): 858–61.
- [27] Kugelberg CF.: Periodontal healing two and four years after impacted lower third molar surgery. A comparative retrospective study. *International Journal of Oral and Maxillofacial Surgery* 1990; 19(6): 341–5.